arXiv:1704.00379v2 [cs.DM] 15 Apr 2017

On the thinness and proper thinness of a graph

Flavia Bonomo[1]

Universidad de Buenos Aires. Facultad de Ciencias Exactas y Naturales. Departamento de Computación. Buenos Aires, Argentina. / CONICET-Universidad de Buenos Aires. Instituto de Investigación en Ciencias de la Computación (ICC). Buenos Aires, Argentina.

Diego de Estrada

Universidad de Buenos Aires. Facultad de Ciencias Exactas y Naturales. Departamento de Computación. Buenos Aires, Argentina.

Abstract

Graphs with bounded thinness were defined in 2007 as a generalization of interval graphs. In this paper we introduce the concept of proper thinness, such that graphs with bounded proper thinness generalize proper interval graphs. We study the complexity of problems related to the computation of these parameters, describe the behavior of the thinness and proper thinness under three graph operations, and relate thinness and proper thinness with other graph invariants in the literature. Finally, we describe a wide family of problems that can be solved in polynomial time for graphs with bounded thinness, generalizing for example list matrix partition problems with bounded size matrix, and enlarge this family of problems for graphs with bounded proper thinness, including domination problems.

Keywords: interval graphs, proper interval graphs, proper thinness, thinness.

Email addresses: fbonomo@dc.uba.ar (Flavia Bonomo), destrada@dc.uba.ar (Diego de Estrada)

[1]**For Pavol Hell on the Occasion of his 70th Birthday**

1. Introduction

A graph $G = (V, E)$ is *k-thin* if there exist an ordering $v_1, \ldots, v_n$ of V and a partition of V into k classes $(V^1, \ldots, V^k)$ such that, for each triple (r, s, t) with $r < s < t$, if v_r, v_s belong to the same class and $v_t v_r \in E$, then $v_t v_s \in E$. The minimum k such that G is k-thin is called the *thinness* of G. The thinness is unbounded on the class of all graphs, and graphs with bounded thinness were introduced in [27] as a generalization of interval graphs, which are exactly the 1-thin graphs. When a representation of the graph as a k-thin graph is given, for a constant value k, some NP-complete problems as maximum weighted independent set and bounded coloring with fixed number of colors can be solved in polynomial time [27, 5]. These algorithms were respectively applied for improving heuristics of two real-world problems: the Frequency Assignment Problem in GSM networks [27], and the Double Traveling Salesman Problem with Multiple Stacks [5]. In this work we propose a framework to describe a wide family of problems that can be solved by dynamic programming techniques on graphs with bounded thinness, when the k-thin representation of the graph is given. These problems generalize for example the list matrix partition problems for matrices of bounded size [14].

We introduce here the concept of *proper thinness* of a graph with the aim of generalizing proper-interval graphs: graphs that are proper 1-thin are exactly proper interval graphs (see Section 3 for a definition). We extend the framework in order to solve in polynomial time by dynamic programming many of the domination-type problems in the literature (e.g. classified in [1]) and their weighted versions, such as existence/minimum (weighted) independent dominating set, minimum (weighted) total dominating set, minimum perfect dominating set and existence/minimum (weighted) efficient dominating set, for the class of graphs with bounded proper thinness k, when the proper k-thin representation of the graph is given.

The organization of the paper is the following. In Section 2 we state the main definitions and present some basic results on thinness. In Sections 2.1 and 3.2, we study some problems related with the recognition of k-thin graphs and proper k-thin graphs. We analyze the computational complexity of finding a suitable vertex partition when a vertex ordering is given, and, conversely, finding a vertex ordering when a vertex partition is given. In Section 2.2 we survey the relation of thinness and other width parameters in graphs.

In Section 3.3 we describe a wide family of problems that can be solved in polynomial time for graphs with bounded thinness, when the representation is given.

In Section 3 we deal with the concept of proper thinness. In Section 3.1 we relate the proper thinness of interval graphs to other interval graph invariants, as interval count and chains of nested intervals. In Section 3.3 we describe a family of dominating-like problems that can be solved in polynomial time for graphs with bounded proper thinness, extending the family already solvable for graphs with bounded thinness.

In Section 4 we describe the behavior of the thinness and proper thinness under three graph operations: union, join, and Cartesian product. The first two results allow us to fully characterize k-thin graphs by forbidden induced subgraphs within the class of cographs. The third result is used to show the polynomiality of the t-rainbow domination problem for fixed t on graphs with bounded proper thinness.

2. Definitions and basic results

All graphs in this work are finite, undirected, and have no loops or multiple edges. For all graph-theoretic notions and notation not defined here, we refer to West [41]. Let G be a graph. Denote by $V(G)$ its vertex set, by $E(G)$ its edge set, by $\overline{G}$ its complement, by $N(v)$ the neighborhood of a vertex v in G, by $N[v]$ the closed neighborhood $N(v) \cup \{v\}$, and by $\overline{N}(v)$ the non-neighbors of v. If $X \subseteq V(G)$, denote by $N(X)$ the set of vertices not in X having at least one neighbor in X.

Denote by $G[W]$ the subgraph of G induced by $W \subseteq V(G)$, and by $G-W$ or $G \setminus W$ the graph $G[V(G) \setminus W]$. A subgraph H (not necessarily induced) of G is a *spanning subgraph* if $V(H) = V(G)$.

Denote the size of a set S by $|S|$. A *clique* (resp. *stable set*) is a set of pairwise adjacent (resp. nonadjacent) vertices. We use *maximum* to mean maximum-sized, whereas *maximal* means inclusion-wise maximal. The use of *minimum* and *minimal* is analogous.

Denote by K_n the graph induced by a clique of size n. A *claw* is the graph isomorphic to $K_{1,3}$. Let H be a graph and t a natural number. The disjoint union of t copies of the graph H is denoted by tH.

For a positive integer r, the $(r \times r)$-*grid* is the graph whose vertex set is $\{(i,j) : 1 \leq i,j \leq r\}$ and whose edge set is $\{(i,j)(k,l) : |i-k| + |j-l| = 1, \text{ where } 1 \leq i,j,k,l \leq r\}$.

A *dominating set* in a graph is a set of vertices such that each vertex outside the set has at least one neighbor in the set.

A *coloring* of a graph is an assignment of colors to its vertices such that any two adjacent vertices are assigned different colors. The smallest number t such that G admits a coloring with t colors (a *t-coloring*) is called the *chromatic number* of G and is denoted by $\chi(G)$. A coloring defines a partition of the vertices of the graph into stable sets, called *color classes*. List variations of the vertex coloring problem can be found in the literature. For a survey on that kind of related problems, see [39]. In the *list-coloring problem*, every vertex v comes equipped with a list of permitted colors $L(v)$ for it.

For a symmetric matrix M over $0, 1, *$, the *M-partition problem* seeks a partition of the vertices of the input graph into independent sets, cliques, or arbitrary sets, with certain pairs of sets being required to have no edges, or to have all edges joining them, as encoded in the matrix M: $M_{ii} = 1$ means the i-th set is a clique, while $M_{ii} = 0$ means the i-th set is a stable set; for $i \neq j$, $M_{ij} = 1$ means every vertex of the i-th set is adjacent to every vertex of the j-th set, while $M_{ij} = 0$ means there are no edges from the i-th set to the j-th set. Moreover, the vertices of the input graph can be equipped with lists, restricting the parts to which a vertex can be placed. In that case the problem is know as a *list matrix partition problem*. Such (list) matrix partition problems generalize (list) colorings and (list) homomorphisms [14].

When discussing about algorithms and data structures, we denote by n the number of vertices of the input graph G.

Given a graph G, a weight function w on $V(G)$, and a subset $S \subseteq V(G)$, the weight of S, denoted by $w(S)$ is defined as $\sum_{v \in S} w(v)$.

A class of graphs is *hereditary* when if a graph G is in the class, then every induced subgraph of G is in the same class.

A graph is a *cograph* if it contains no induced path of length four.

A graph $G(V, E)$ is a *comparability graph* if there exists an ordering $v_1, \ldots, v_n$ of V such that, for each triple (r, s, t) with $r < s < t$, if $v_r v_s$ and $v_s v_t$ are edges of G, then so is $v_r v_t$. Such an ordering is a *comparability ordering*. A graph is a *co-comparability graph* if its complement is a comparability graph.

An graph G is an *interval graph* if to each vertex $v \in V(G)$ a closed interval $I_v = [l_v, r_v]$ of the real line can be associated, such that two distinct vertices $u, v \in V(G)$ are adjacent if and only if $I_u \cap I_v \neq \emptyset$. The family $\{I_v\}_{v \in V(G)}$ is an *interval representation* of G. An undirected graph G is a *proper interval graph* if there is an interval representation of G in which no

interval properly contains another. In the same way, an undirected graph G is a *unit interval graph* if there is an interval representation of G in which all the intervals have the same length.

In 1969, Roberts [32] proved that the classes of proper interval graphs, unit interval graphs, and interval graphs with no claw as induced subgraph coincide.

The right-end ordering of the vertices of an interval graph satisfies the following property: for each triple (r, s, t) with $r < s < t$, if $v_t v_r \in E$, then $v_t v_s \in E$. In other words, the neighbors of vertex t with index less than t are $t-1, t-2, \ldots, t-d$ for some $d \geq 0$. Moreover, a graph G is an interval graph if and only if there exists an ordering of its vertices satisfying the property above [31, 29].

Let us repeat and extend the definition of k-thinness given in the introduction. A graph $G = (V, E)$ is *k-thin* if there exist an ordering $v_1, \ldots, v_n$ of V and a partition of V into k classes such that, for each triple (r, s, t) with $r < s < t$, if v_r, v_s belong to the same class and $v_t v_r \in E$, then $v_t v_s \in E$. An ordering and a partition satisfying those properties are said to be *consistent*. The minimum k such that G is k-thin is called the *thinness* of G and denoted by $\text{thin}(G)$.

The *thinness* of a graph was introduced by Mannino, Oriolo, Ricci, and Chandran in 2007 [27]. Graphs with *bounded thinness* (thinness bounded by a constant value) are a generalization of interval graphs, that are exactly the graphs with thinness 1, and capture some of their algorithmic properties.

Let $\overline{tK_2}$ be the complement of a matching of size t.

Theorem 1. [27] *For every $t \geq 1$, $\text{thin}(\overline{tK_2}) = t$.*

2.1. Algorithmic aspects

We will deal in this section with some questions related to the recognition problem of k-thin graphs. The recognition problem itself is open so far, but we will show that, given a vertex ordering of a graph, we can find in polynomial time a partition into a minimum number of classes which is consistent with the ordering. On the other hand, we will show that given a graph and a vertex partition, it is NP-complete to decide if there exists an ordering of the vertices of the graph which is consistent with the partition.

Theorem 2. *Given a graph G and an ordering $<$ of its vertices, one can find in polynomial time a graph $G_<$ with the following properties:*

(1) $V(G_<) = V(G)$*;*

(2) the chromatic number of $G_<$ is equal to the minimum integer k such that there is a partition of $V(G)$ into k sets that is consistent with the order $<$, and the color classes of a valid coloring of $G_<$ form a partition consistent with $<$;

(3) $G_<$ is a co-comparability graph.

In particular, the minimum integer k as in (2) and a partition into k vertex sets can be computed in polynomial time. Moreover, if G is a co-comparability graph and $<$ a comparability ordering of $\overline{G}$, then $G_<$ is a spanning subgraph of G.

Proof. Let G be a graph and $<$ an ordering of its vertices. We will build a graph $G_<$ such that $V(G_<) = V(G)$, and $v < w$ are adjacent in $G_<$ if and only if they cannot belong to the same class of a partition which is consistent with $<$. By definition of consistence, this happens if and only if there is a vertex z in G such that $v < w < z$, z is adjacent to v and nonadjacent to w. So define $E(G_<)$ such that for $v < w$, $vw \in E(G_<)$ if and only if there is a vertex z in G such that $v < w < z$, $zv \in E(G)$ and $zw \notin E(G)$.

Let us see that $<$ is a comparability ordering for $\overline{G_<}$. Suppose on the contrary that there is a triple $r < s < t$ in $V(G)$ such that rs, st are edges of $\overline{G_<}$ and rt is not an edge of $\overline{G_<}$. By definition of $G_<$, there is a vertex z such that $r < s < t < z$, $zr \in E(G)$ and $zt \notin E(G)$. If $zs \notin E(G)$, then rs is an edge of $G_<$, a contradiction. If $zs \in E(G)$, then st is an edge of $G_<$, a contradiction as well. So $G_<$ is a co-comparability graph, being $<$ a comparability ordering for $\overline{G_<}$.

As we have defined $G_<$ such that $V(G_<) = V(G)$, and $v < w$ are adjacent in $G_<$ if and only if they cannot belong to the same class of a partition which is consistent with $<$, it follows that there is a one-to-one correspondence between partitions of $V(G)$ consistent with $<$ and colorings of $G_<$. In particular, the minimum k such that there is a partition of $V(G)$ into k sets that is consistent with $<$ is the chromatic number of $G_<$. An optimum coloring of $G_<$ can be computed in polynomial time [16].

To complete the proof of the theorem, suppose now that G is a co-comparability graph and $<$ is a comparability ordering for $\overline{G}$. Let $v < w$

adjacent in $G_<$. By definition, there is a vertex z in G such that $v < w < z$, $vz \in E(G)$ and $wz \notin E(G)$. If $vw \notin E(G)$, being $\overline{G}$ a comparability graph, $vz \notin E(G)$, a contradiction. So $vw \in E(G)$. This proves that $G_<$ is a spanning subgraph of G. □

A direct consequence of this result is the following.

Corollary 3. [5] *If G is a co-comparability graph,* $\text{thin}(G) \leq \chi(G)$. *Moreover, any vertex partition given by a coloring of G and any comparability ordering for its complement are consistent.*

As already observed in [5], the bound $\text{thin}(G) \leq \chi(G)$ for co-comparability graphs can be arbitrarily bad: for example, if G is a clique of size n, then $\text{thin}(G) = 1$ and $\chi(G) = n$. However, it holds with equality for graphs $\overline{tK_2}$, because $\text{thin}(\overline{tK_2}) = \chi(\overline{tK_2}) = t$ (Theorem 1).

In contrast with Theorem 2, if a partition is given, it is NP-complete to decide the existence of a consistent ordering.

Consistent ordering with a given partition
Instance: A graph $G = (V, E)$ and a partition of V into non-empty subsets.
Question: Does there exist a total order $<$ of V consistent with the given partition?

The proof is based on a reduction from the following problem, which is known to be NP-complete [18].

Non-Betweenness
Instance: A finite set A and a collection of ordered triples S of elements of A.
Question: Does there exist a total order $<$ of A such that for each $(x, y, z) \in S$, it is never the case that $x < y < z$ or $z < y < x$ (i.e. y is *not between* x and z)?

We start with an easy lemma.

Lemma 4. *Let G be a graph, $<$ an ordering of $V(G)$ and $V_1, \ldots, V_k$ a partition of $V(G)$ that is consistent with $<$. Let $\{x_i, y_i\} \subseteq V_i$, for $i = 1, 2$, such*

that x_1x_2 and y_1y_2 are the only edges between $\{x_1, y_1\}$ and $\{x_2, y_2\}$. Then $x_1 < y_1$ if and only if $x_2 < y_2$.

Proof. By symmetry, let us assume that y_1 is the biggest vertex according to $<$. Again by symmetry, to prove the lemma it is enough to prove that $x_2 < y_2$. But by definition of consistence, since x_2 and y_2 are in the same class and y_1 is adjacent to y_2 but not to x_2, it is not possible that $y_2 < x_2 < y_1$. □

Theorem 5. *The problem* Consistent ordering with a given partition *is NP-complete.*

Proof. First note that Consistent ordering with a given partition is in NP, by using the total order of V as the certificate.

Now let us prove its NP-hardness. Given an instance (A, S) of Non-Betweenness, build a graph $G = (V, E)$ and a partition $V_0, V_1, \ldots V_{|S|}$ of V as follows.

Fix an ordering of the triples in S. Nodes of V_0 are in one-to-one correspondence with elements of A. For $i = 1, \ldots, |S|$, V_i has 3 nodes, and they are in a one-to-one correspondance with the elements of the i-th triple in S. Let us call a^i the element of V_i that corresponds to $a \in A$, for $i = 0, \ldots, |S|$.

Define the edges of G as follows: for each triple $(x, y, z) \in S$, let V_i be its corresponding set. The only edge in the subgraph induced by $\{x^i, y^i, z^i\}$ is x^iz^i. The remaining edges of G are all the possible edges between vertices associated to the same $a \in A$.

Suppose first there is an ordering $<$ consistent with the partition $\{V_0, \ldots, V_{|S|}\}$. By Lemma 4, for each $1 \leq i \leq |S|$, the relative order of the vertices x^i, y^i, z^i is the same as the relative order of the vertices x^0, y^0, z^0. By definition of consistence and since the only edge in the subgraph induced by $\{x^i, y^i, z^i\}$ is x^iz^i, y^i is not between x^i and z^i in that order. So the order of the vertices in V_0 gives a positive answer to the instance (A, S) of Non-Betweenness.

Suppose now that there is a valid order $<$ for the instance (A, S) of Non-Betweenness. We can extend $<$ to $V(G)$ by making consecutive all the copies in $V(G)$ of an element of A. Now, let $p < q < r$ be three vertices of G such that p, q belong to the same class V_i and $rp \in E(G)$. Since V_0 is a stable set and the triples in S satisfy the non-betweenness condition, r is not in V_i. So r and p correspond to the same element a of A, and since

there is at most one copy of an element of A in each V_i, q does not correspond to a copy of a. But this contradicts the fact that all the vertices of G that correspond to a same element of A are consecutive. So the situation cannot arise, and the extended order is consistent with the partition. □

The computational complexity of the decision of existence of a consistent ordering when the number of sets in the partition is fixed is still open. So is the computational complexity of deciding if a graph is k-thin, even for fixed $k \geq 2$.

2.2. Thinness and other width parameters

Many width parameters are defined in the literature. In this section we compile the results relating the thinness with some of them, namely pathwidth [34], treewidth [2, 35], clique-width [10], and boxicity [33].

A *path decomposition* of a graph $G = (V, E)$ is a sequence of subsets of vertices $(X_1, \ldots, X_r)$ such that

1. $X_1 \cup \cdots \cup X_r = V$.
2. for all edges $vw \in E$, there exists an i, $1 \leq i \leq r$, with $\{v, w\} \subseteq X_i$.
3. for all $1 \leq i \leq j \leq k \leq r$, $X_i \cap X_k \subseteq X_j$.

The *width* of the path decomposition is defined as $\max_{1 \leq i \leq r} |X_i| - 1$. The *pathwidth* of a graph G is the minimum width over all path decompositions of G.

A *tree decomposition* of a graph $G = (V_G, E_G)$ is defined by a rooted tree $T = (V_T, E_T)$ and a mapping of $v \in V_G$ to a nonempty subset of V_T such that, if X_z is the set of vertices of G mapped to vertex $z \in V_T$,

1. for all edges $vw \in E_G$, there exists $z \in V_T$, with $\{v, w\} \subseteq X_z$.
2. for every vertex $v \in V_G$, the subset of V_T to which it is mapped is a connected subgraph of T.

The *width* of the tree decomposition is defined as $\max_{z \in V_T} |X_z| - 1$. The *treewidth* of a graph G is the minimum width over all tree decompositions of G.

The *clique-width* of a graph G is defined as the minimum number of labels needed to construct G by means of the following operations:

1. Creation of a new vertex v with label i.
2. Disjoint union of two labeled graphs G and H.
3. Joining by an edge every vertex labeled i to every vertex labeled j, where $i \neq j$.
4. Renaming label i to label j.

For a graph G, the *boxicity* is defined as the minimum dimension d such that G is the intersection graph of boxes with edges parallel to the axes in d-dimensional space. In particular, a graph has boxicity 2 if and only if it is the intersection graph of rectangles with edges parallel to the axes. Graphs with boxicity one are exactly the interval graphs.

In [27] it was proved that the thinness of a graph is at most the pathwidth plus one, and that the gap may be high, since the pathwidth of a complete graph with r vertices is $r - 1$, while its thinness is 1.

On the other hand, in [8] it was proved that the boxicity is a lower bound for the thinness of a graph, and it was pointed out that the difference can be large, as an $(r \times r)$-grid has boxicity 2 and thinness $\Theta(r)$.

The *vertex isoperimetric peak* of a graph G, denoted as $b_v(G)$, is defined as $\max_s \min_{X \subset V, |X|=s} |N(X)|$. The thinness of the grid was estimated by using the following result, that was also used in [3] to give a lower bound of the thinness of a complete binary tree. We will use use it as well to estimate the thinness of complete m-ary trees.

Lemma 6. [8] *For every graph G,* $\text{thin}(G) \geq b_v(G)/\Delta(G)$.

Interval graphs have thinness 1 and unbounded clique-width [17], while cographs have clique-width 2 [11] and unbounded thinness, because $\overline{tK_2}$ is a cograph for every t, so the parameters are not comparable.

Complete graphs have high treewidth and thinness 1, and trees instead have treewidth 1 but we have the following result.

Theorem 7. *For every fixed value m, the thinness of the complete m-ary tree on n vertices is $\Theta(\log n)$.*

Proof. In [40] it was proved that the vertex isoperimetric peak of the complete m-ary tree of height h is $\Theta(h)$. On the other hand, it was proved in [13, 37] that the pathwidth of the complete m-ary tree of height h is $\Theta(h)$. As the thinness of a graph is upper bounded by the pathwidth plus one [27] and using Lemma 6, it follows that the thinness of the complete m-ary tree of height h is $\Theta(h)$, and this proves the theorem. □

2.3. Solving combinatorial optimization problems on graphs with bounded thinness

Since a k-thin graph G does not contain $\overline{(k+1)K_2}$ as induced subgraph (Theorem 1), it has at most $|V(G)|^{2k}$ maximal cliques [30]. In particular, the maximum weighted clique problem can be solved in polynomial time on graphs with bounded thinness, by simple enumeration of the maximal cliques of the graph [38].

The maximum weighted stable set problem can be solved in polynomial time on graphs with bounded thinness, when an ordering and a partition that are consistent are given [27]. In the same hypothesis, the capacitated coloring (in which there is an upper bound α_j on the number of vertices of color j) can be solved in polynomial time, if the number of colors s is fixed [5]. As a byproduct, in the same paper it is shown that the capacitated coloring can be solved in polynomial time for co-comparability graphs, if the number of colors s is fixed, in contrast with the case in which the bounds α_j are all equal to a fixed number h, that is NP-complete, even for two subclasses of co-comparability graphs: permutation graphs (for $h \geq 6$) [25] and interval graphs (for $h \geq 4$) [4]. The hardness on interval graphs implies the hardness for graphs of bounded thinness, since interval graphs are the graphs with thinness 1.

Both algorithms, the one for maximum weighted stable set and the one for capacitated coloring with fixed number of colors, are based on dynamic programming. One of the main results in this work is a generalization of these algorithmic results. We describe now a framework of problems that can be solved for graphs with bounded thinness, given the representation.

Instance:

- A k-thin representation of $G = (V, E)$, with ordering $<$ of V, namely $v_1 < \cdots < v_n$, and partition of V into k classes $V^1, \ldots, V^k$.
- A family of arbitrary nonnegative weights $w_1, \ldots, w_t$ on V.
- A family of nonnegative weights $b_1, \ldots, b_p$ on V bounded by a fixed polynomial in n (p fixed, $q(n)$ the bound for the weights).

Question: find sets $S_1, \ldots, S_r$ (r fixed, not necessarily disjoint), $S_j \subseteq V$ for $1 \leq j \leq r$, such that:

- the objective is to minimize or maximize a linear function $\sum_{1 \leq i \leq t; 1 \leq j \leq r} c_{ij} w_i(S_j)$.
- each vertex v has a list $L(v)$ of combinations of the sets $S_1, \ldots, S_r$ to which it can belong (that may include the empty combination).
- there is an $r \times r$ symmetric matrix M over $0, 1, *$, stating the adjacency conditions on the sets S_j, such that for $1 \leq i < j \leq r$, $M_{ii} = 1$ means S_i is a clique, $M_{ii} = 0$ means S_i is a stable set, $M_{ij} = 1$ means all the edges joining S_i and S_j have to be present, $M_{ij} = 0$ means there are no edges from S_i to S_j.
- there is a family of restrictions of type
  - $0 \leq l_{iJ\cap} \leq b_i(\bigcap_{j \in J} S_j) \leq u_{iJ\cap}$, such that $1 \leq i \leq p$, $J \subseteq \{1, \ldots, r\}$.
  - $0 \leq l_{iJ\cup} \leq b_i(\bigcup_{j \in J} S_j) \leq u_{iJ\cup}$, such that $1 \leq i \leq p$, $J \subseteq \{1, \ldots, r\}$.

The family of problems that can be modeled within this framework includes weighted variations of list matrix partition problems with matrices of bounded size, which in turn generalize coloring, list coloring, list homomorphism, equitable coloring with different objective functions, all for fixed number of colors (or graph size in the case of homomorphism), clique cover with fixed number of cliques, weighted stable sets, and other graph partition problems. It models also sum-coloring and its more general version optimum cost chromatic partition problem [22] for fixed number of colors, but it does

not include dominating-like problems.

We will solve such a problem as a shortest or longest path problem (according to minimization or maximization of the objective function) in an auxiliary acyclic digraph $D = (X, A)$ whose nodes correspond to *states* and whose arcs are weighted and labeled. The total weight of the path is the value of the objective function in the solution that can be builded by using the arc labels.

A state is a tuple, containing:

- a number $1 \leq s \leq |V_G|$ indicating that we are considering the subgraph G_s of G, induced by $v_1, \ldots, v_s$.
- nonnegative parameters $l_{iJ\cap}, u_{iJ\cap}, l_{iJ\cup}, u_{iJ\cup}$, for $1 \leq i \leq p$, $J \subseteq \{1, \ldots, r\}$; they are at most $2^{r+2}p$, and each of them may take a nonnegative value at most $nq(n)$, which is an upper bound for $b_i(V)$, for every $1 \leq i \leq p$.
- a family of nonnegative parameters $\{\alpha_{ij}\}_{1 \leq i \leq k; 1 \leq j \leq r}$, meaning that we cannot pick for S_j a vertex of the first α_{ij} vertices of the set V^i of the partition; they are kr and each of them may take a nonnegative value at most $n - 1$.
- a family of nonnegative parameters $\{\beta_{ij}\}_{1 \leq i \leq k; 1 \leq j \leq r}$, meaning that we cannot pick for S_j a vertex on the last β_{ij} vertices of the set V^i of the partition; they are kr and each of them may take a nonnegative value at most $n - 1$.

The total number of states is then at most $n^{2kr+1}(nq(n))^{2^{r+2}p}$, that is polynomial in n, since k, r, and p are constant and $q(n)$ is polynomial in n.

The digraph D will have nodes that correspond to possible states, organized in layers $X_0, X_1, \ldots, X_n$ such that X_0 contains only one node x_0, and the layer X_s contains the states whose first parameter is s. The layer X_n contains also only one node, corresponding to the state $(n, \{l_{iJ\cap}\}, \{u_{iJ\cap}\}, \{l_{iJ\cup}\}, \{u_{iJ\cup}\}, \{\alpha_{ij}\}, \{\beta_{ij}\})$, where the parameters $\{l_{iJ\cap}\}, \{u_{iJ\cap}\}, \{l_{iJ\cup}\}, \{u_{iJ\cup}\}$ are the ones in the original formulation of the problem and $\alpha_{ij} = \beta_{ij} = 0$ for every $1 \leq i \leq k$, $1 \leq j \leq r$.

All arcs of A have the form (u, w) with $u \in X_s$ and $w \in X_{s+1}$, for some $0 \leq r \leq n - 1$.

We associate with each node of X a suitable problem, in the same framework, whose parameters correspond to the parameters in the state, but with additional constraints associated with the parameters $\{\alpha_{ij}\}$ and $\{\beta_{ij}\}$.

We will define the arcs in such a way that a node is reachable from the node in the layer X_0 if and only if the associated problem has a solution. The length of the path will be the weight of the solution, and the set of arc labels will encode the solution. Let us describe the arcs of the digraph.

Let w be a node with parameters $(1, \{l_{iJ\cap}\}, \{u_{iJ\cap}\}, \{l_{iJ\cup}\}, \{u_{iJ\cup}\}, \{\alpha_{ij}\}, \{\beta_{ij}\})$.

Let $1 \leq \ell \leq k$ such that $v_1 \in V^\ell$. For each $\tilde{J} \in L(v_1)$ (in particular $\tilde{J} \subseteq \{1, \ldots, r\}$), such that:

1.1 For each $j \in \tilde{J}$, $\beta_{\ell j} = \alpha_{\ell j} = 0$.

1.2 For each $J \subseteq \tilde{J}$, $l_{iJ\cap} \leq b_i(v_1) \leq u_{iJ\cap}$.

1.3 For each $J \not\subseteq \tilde{J}$, $l_{iJ\cap} = 0$.

1.4 For each J such that $J \cap \tilde{J} \neq \emptyset$, $l_{iJ\cup} \leq b_i(v_1) \leq u_{iJ\cup}$.

1.5 For each J such that $J \cap \tilde{J} = \emptyset$, $l_{iJ\cup} = 0$.

we add an arc from x_0 to w, labeled by $\tilde{J}$ and of weight $\sum_{1 \leq i \leq t; j \in \tilde{J}} c_{ij} w_i(v_1)$. If no $\tilde{J}$ satisfies conditions 1.1–1.5, no arc ending in w is added. If more than one arc $x_0 w$ was added, we can keep only the one with maximum (resp. minimum) weight if we are solving a maximization (resp. minimization) problem.

Note that if we add the arc $x_0 w$ labeled by $\tilde{J}$, then the solution $S_j = \{v_1\}$ for $j \in \tilde{J}$, $S_j = \emptyset$ for $j \notin \tilde{J}$ has weight $\sum_{1 \leq i \leq t; j \in \tilde{J}} c_{ij} w_i(v_1)$ and satisfies the state described by w: condition 1.1 says that v_1 (the first and last vertex of V^ℓ in G_1) is allowed to be picked for every set S_j for $j \in \tilde{J}$; conditions 1.2–1.5 say that the assignment does not violate weight constraints.

Let w be a node with parameters $(s, \{l_{iJ\cap}\}, \{u_{iJ\cap}\}, \{l_{iJ\cup}\}, \{u_{iJ\cup}\}, \{\alpha_{ij}\}, \{\beta_{ij}\})$, $1 < s \leq n$.

Let $1 \leq \ell \leq k$ such that $v_s \in V^\ell$. For each $\tilde{J} \in L(v_s)$, such that:

s.1 For each $j \in \tilde{J}$, $\beta_{\ell j} = 0$.

s.2 For each $j \in \tilde{J}$, $\alpha_{\ell j} < |V^\ell \cap \{v_1, \ldots, v_s\}|$.

$s.3$ For each $J \subseteq \tilde{J}$, $b_i(v_s) \leq u_{iJ\cap}$.

$s.4$ For each J such that $J \cap \tilde{J} \neq \emptyset$, $b_i(v_s) \leq u_{iJ\cup}$.

we add an arc from u to w, labeled by $\tilde{J}$ and of weight $\sum_{1\leq i\leq t; j\in\tilde{J}} c_{ij}w_i(v_s)$, where u has parameters $(s-1,\ \{l'_{iJ\cap}\},\ \{u'_{iJ\cap}\},\ \{l'_{iJ\cup}\},\ \{u'_{iJ\cup}\},\ \{\alpha'_{ij}\},\ \{\beta'_{ij}\})$, such that:

$s'.1$ Let $1 \leq j \leq r$. If there exists $j' \in \tilde{J}$ such that $M_{jj'} = 0$, then $\beta'_{\ell j} = \max\{\beta_{\ell j}-1, |N(v_s)\cap V^\ell \cap \{1,\ldots,s-1\}|\}$, and for $1 \leq i \leq k$, $i \neq \ell$, $\beta'_{ij} = \max\{\beta_{ij}, |N(v_s)\cap V^i \cap \{1,\ldots,s-1\}|\}$. Otherwise, $\beta'_{\ell j} = \max\{0, \beta_{\ell j}-1\}$, and for $1 \leq i \leq k$, $i \neq \ell$, $\beta'_{ij} = \beta_{ij}$.

$s'.2$ Let $1 \leq j \leq r$. If there exists $j' \in \tilde{J}$ such that $M_{jj'} = 1$, then $\alpha'_{\ell j} = \max\{\min\{|V^\ell \cap \{1,\ldots,s-1\}|, \alpha_{\ell j}\}, |\overline{N}(v_s)\cap V^\ell \cap \{1,\ldots,s-1\}|\}$, and for $1 \leq i \leq k$, $i \neq \ell$, $\alpha'_{ij} = \max\{\alpha_{ij}, |\overline{N}(v_s) \cap V^i \cap \{1,\ldots,s-1\}|\}$. Otherwise, $\alpha'_{\ell j} = \min\{|V^\ell \cap \{1,\ldots,s-1\}|, \alpha_{\ell j}\}$, and for $1 \leq i \leq k$, $i \neq \ell$, $\alpha'_{ij} = \alpha_{ij}$.

$s'.3$ For each $J \subseteq \tilde{J}$, $l'_{iJ\cap} = \max\{0, l_{iJ\cap} - b_i(v_s)\}$ and $u'_{iJ\cap} = u_{iJ\cap} - b_i(v_s)$.

$s'.4$ For each $J \not\subseteq \tilde{J}$, $l'_{iJ\cap} = l_{iJ\cap}$ and $u'_{iJ\cap} = u_{iJ\cap}$.

$s'.5$ For each J such that $J \cap \tilde{J} \neq \emptyset$, $l'_{iJ\cup} = \max\{0, l_{iJ\cup} - b_i(v_s)\}$ and $u'_{iJ\cup} = u_{iJ\cup} - b_i(v_s)$.

$s'.6$ For each J such that $J \cup \tilde{J} = \emptyset$, $l'_{iJ\cup} = l_{iJ\cup}$ and $u'_{iJ\cup} = u_{iJ\cup}$.

If no $\tilde{J}$ satisfies conditions $s.1$–$s.4$, no arc ending in w is added. If more than one arc from the same vertex u to w was added, we can keep only the one with maximum (resp. minimum) weight if we are solving a maximization (resp. minimization) problem.

That is, if an arc is added, the arc corresponds to the choice of adding the vertex v_s to the sets $\{S_j\}_{j\in\tilde{J}}$, the conditions required imply that the choice is valid for w in the case that the state described by u admits a solution, the label of the arc keeps track of the choice made, and the cost corresponds to the weight that the choice adds to the objective function.

Note that if we add the arc uw labeled by $\tilde{J}$, then for a solution $\{S'_j\}_{1\leq j\leq r}$ for G_{s-1} satisfying the state described by u, then the solution $\{S_j\}_{1\leq j\leq r}$ for

G_s such that $S_j = S'_j \cup \{v_s\}$ for $j \in \tilde{J}$, $S_j = S'_j$ for $j \notin \tilde{J}$ satisfies the state described by w. Conditions $s.1$ and $s.2$ say that v_s (the last vertex of V^ℓ in G_s) is allowed to be picked for every set S_j for $j \in \tilde{J}$. Condition $s'.1$ ensures on one hand that the conditions imposed by the parameters $\{\beta_{ij}\}$ in w are satisfied by the solution of u, and, on the other hand, that if $j' \in \tilde{J}$ and $1 \leq j \leq r$ are such that $M_{jj'} = 0$ then no neighbor of v_s belongs to S'_j, as required. Similarly, condition $s'.2$ ensures on one hand that the conditions imposed by the parameters $\{\alpha_{ij}\}$ in w are satisfied also by the solution of u, and, on the other hand, that if $j' \in \tilde{J}$ and $1 \leq j \leq r$ are such that $M_{jj'} = 1$ then all vertices in S'_j are adjacent to v_s, as required. These conditions strongly use that the order and the partition are consistent. Finally, conditions $s.3$–$s.4$, and $s'.3$–$s'.6$ ensure that the solution does not violate weight constraints.

Moreover, the difference of weight of the solution $\{S_j\}_{1 \leq j \leq r}$ with respect to $\{S'_j\}_{1 \leq j \leq r}$ is exactly $\sum_{1 \leq i \leq t; j \in \tilde{J}} c_{ij} w_i(v_s)$.

In that way, a directed path in the digraph corresponds to an assignment of vertices of the graph to lists of sets and its weight is the value of the objective function for the corresponding assignment.

The digraph has a polynomial number of nodes and can be builded in polynomial time. Since it is acyclic, both the longest path and shortest path can be computed in linear time in the size of the digraph by topological sorting.

Remark 1. *A similar algorithm for the same family of problems can be described for co-k-thin graphs (complements of k-thin graphs), k bounded.*

3. Proper thinness

The right-end ordering of the vertices of a proper interval graph satisfies the following property: for each triple (r, s, t) with $r < s < t$, if $v_t v_r \in E$, then $v_t v_s \in E$ and $v_r v_s \in E$. In other words, the neighbors of vertex t with index less than t are $t-1, t-2, \ldots, t-d$, and those with index greater than t are $t+1, t+2, \ldots, t+d'$. Moreover, G is a proper interval graph if and only if there exists an ordering of its vertices satisfying the property above [12, 26].

We define the concept of *proper thinness* as follows.

A graph $G = (V, E)$ is *proper k-thin* if there exist an ordering $v_1, \ldots, v_n$ of V and a partition of V into k classes $(V^1, \ldots, V^k)$ such that, for each

triple (r, s, t) with $r < s < t$, if v_r, v_s belong to the same class and $v_t v_r \in E$, then $v_t v_s \in E$ and if v_s, v_t belong to the same class and $v_r v_t \in E$, then $v_r v_s \in E$. Equivalently, G is proper k-thin if both $v_1, \ldots, v_n$ and $v_n, \ldots, v_1$ are consistent with the partition. In this case, the partition and the ordering $v_1, \ldots, v_n$ are said to be *strongly consistent*, and the minimum k such that G is proper k-thin is called the *proper thinness* of G and denoted by $\text{pthin}(G)$.

Since k-thin graphs are defined as a generalization of interval graphs, proper k-thin graphs arise naturally as a generalization of proper interval graphs. It can be seen that a graph is proper 1-thin if and only if it is a proper interval graph. Moreover, the proper-thinness of the class of interval graphs is unbounded (See Proposition 8).

But the real aim of this definition is to be able to deal with a more general family of problems, including domination-type ones, on graphs with bounded proper thinness. We will give more details in Section 3.3.

3.1. Interval graphs with high proper thinness

The proper thinness of the class of interval graphs is unbounded. A family of interval graphs with arbitrarily large proper thinness is the following.

Let $h \geq 1$, and define $claw_h$ as the graph obtained from the complete ternary tree of height h by adding all the edges between a vertex of the tree and its ancestors. It is easy to see that claw_h is an interval graph for every $h \geq 1$ (an interval representation of claw_3 can be seen in Figure 1). The graph claw_1 is the claw.

Proposition 8. [36] *For any $h \geq 1$, $\text{pthin}(claw_h) = h + 1$.*

Proof. Let $h \geq 1$. We will label the vertices of $G = \text{claw}_h$ as v_j^i such that $0 \leq i \leq h$, $1 \leq j \leq 3^i$, v_1^0 is the root of the ternary tree, and for each $0 \leq i \leq h-1$, $1 \leq j \leq 3^i$, the children of v_j^i are v_{3j-2}^{i+1}, v_{3j-1}^{i+1}, and v_{3j}^{i+1}. Let us consider an ordering $<$ and a partition of $V(G)$ that are strongly consistent. Without loss of generality, by symmetry, we may assume $v_2^i < v_1^i < v_3^i$ for every $i \geq 1$.

Let us show now that for every $0 \leq i' < i \leq h$, v_1^i and $v_1^{i'}$ cannot be in the same class of the partition. Otherwise, if $v_1^i < v_1^{i'}$ then the fact of $v_2^i < v_1^i < v_1^{i'}$, $v_2^i v_1^{i'} \in E(G)$ and $v_2^i v_1^i \notin V(G)$ contradicts the definition of strong consistence, and if $v_1^{i'} < v_1^i$ then the fact of $v_1^{i'} < v_1^i < v_3^i$, $v_3^i v_1^{i'} \in E(G)$ and $v_3^i v_1^i \notin V(G)$ contradicts the definition of strong consistence.



Figure 1: An interval representation of claw_3.

So, $v_1^0, \ldots, v_1^h$ are all in different classes of the partition and $\text{pthin}(\text{claw}_h) \geq h+1$. On the other hand, a partition of the vertices according to its height in the tree, and a postorder of the vertices of the tree are strongly consistent. Thus $\text{pthin}(\text{claw}_h) = h+1$. □

This example is also a classical example of a graph with high interval count and high length of a chain of nested intervals. We will relate the proper thinness of interval graphs with these two interval graphs invariants.

The *interval count* of an interval graph G is the minimum number of different interval sizes needed in an interval representation of G (see for example [7, 24]). Graphs with interval count at most k are also known as *k-length interval graphs*.

A *k-nested interval graph* is an interval graph admitting an interval representation in which there are no chains of $k+1$ intervals nested in each other [23]. It is easy to see that k-nested interval graphs are a superclass of k-length interval graphs. We have also the following property.

Proposition 9. [28] *Every k-nested interval graph is proper k-thin.*

Proof. Let G be a k-nested interval graph and consider an interval representation of G with no chains of $k+1$ intervals nested in each other. It is a known result that we may assume that all the interval endpoints are distinct. We label each interval by the length of the longest chain of nested intervals ending in it, and these labels define the partition of the vertices into classes, that are at most k. Now, we order the vertices according to their intervals by the right endpoint (left to right). That order is consistent with the partition in which the only class contains all vertices of G, so, in particular, it is consistent with every other partition refining it. Let us see that the consistence is strong. Let $r < s < t$ such that s and t are in the same class of the partition. Let I_r, I_s, I_t their corresponding intervals. By definition of the classes, $I_s \not\subseteq I_t$, otherwise the length of the longest chain of nested intervals ending in I_s would be strictly greater than the one for I_t.

As the right endpoint of I_t is greater than the one of I_s, it follows that the left endpoint of I_t is also greater than the one of I_s. Thus, if I_r intersects I_t, it intersects I_s as well. So, the ordering and the partition are strongly consistent and G is proper k-thin. □

Graphs with interval count one are known as unit interval graphs, while 1-nested interval graph are equivalent to proper interval graphs. In [32] it is showed that unit interval graphs are equivalent to proper interval graphs. So the classes proper 1-thin, 1-length interval and 1-nested interval are equivalent. We will see that for higher numbers the equivalence does not necessarily hold.

Indeed, in [15, Theorem 5, p. 177], Fishburn shows that, for every $k \geq 2$, there are 2-nested interval graphs that are not k-length interval.

We will describe a family of graphs that show that, for every $k \geq 3$, there are proper 3-thin graphs that are not k-nested interval.

Let $k \geq 1$. Let G_k with $3k+1$ vertices is defined as follows. Its vertex-set is $V_k = A_k \cup B_k \cup W_k$, where $A_k = \{a_1, \ldots, a_k\}$, $B_k = \{b_1, \ldots, b_k\}$ and $W_k = \{v_1, \ldots, v_k, v_{k+1}\}$. The subgraph induced by W_k is a clique with $k+1$ vertices; a_1 (resp., b_1) is adjacent to v_1. Then, for any $1 < i \leq k$, a_i (resp., b_i) is adjacent to a_{i-1} (resp., to b_{i-1}), and to v_j for any $j \geq i$. See Figure 2 for a sketch of G_k and an interval representation of it.

The graph G_1 is the claw, which is not proper interval. For higher values of k, we have the following property.

Proposition 10. [28] *For any $k \leq 2$, G_k is proper 3-thin, but in every interval representation of it, if I_j is the interval corresponding to v_j, it holds $I_{k+1} \subseteq I_k \subseteq \cdots \subseteq I_1$.*

Proof. Consider the ordering $a_1, \ldots, a_k, b_1, \ldots, b_k, v_1, \ldots, v_k, v_{k+1}$, and the three classes A_k, B_k and W_k. It is easy to see that are strongly consistent.

Let $1 \leq i \leq k-1$. Notice that $a_i a_{i+1} v_{i+1} b_{i+1} b_i$ induce a path of length five on G_k. In every interval representation of it, the interval I_{i+1} is between the intervals corresponding to a_i and b_i and disjoint to them. As the five vertices are adjacent to v_i, it follows that the $I_{i+1} \subseteq I_i$. Finally, by the shape of interval representations of a path of length five, each of the intervals corresponding to a_k and b_k contains an endpoint of I_k. As v_{k+1} is neither adjacent to a_k nor to b_k, $I_{k+1} \subseteq I_k$. □

The following characterization was proved for k-nested interval graphs.



Figure 2: A sketch of graph G_k and an interval representation of it.

Lemma 11. [23] *An interval graph is k-nested interval if and only if it has an interval representation which can be partitioned into k proper interval representations.*

This lemma and the family of graphs G_k show that even if the vertices of a proper k-thin graph can be partitioned into k sets of vertices each of them inducing a proper interval graph, it is not always the case that it has an interval representation which can be partitioned into k proper interval representations.

3.2. Algorithmic aspects

In this section we will show that many of the properties of thinness showed in Section 2.1 hold similarly for proper thinness.

Theorem 12. *Given a graph G and an ordering $<$ of its vertices, one can find in polynomial time a graph $G_<$ with the following properties:*

(1) $V(G_<) = V(G)$*;*

(2) the chromatic number of $G_<$ is equal to the minimum integer k such that there is a partition of $V(G)$ into k sets that is strongly consistent with the order $<$, and the color classes of a valid coloring of $G_<$ form a partition strongly consistent with $<$;

(3) $G_<$ is a co-comparability graph.

In particular, the minimum integer k as in (2) and a partition into k vertex sets can be computed in polynomial time. Moreover, if G is a co-comparability graph and $<$ a comparability ordering of $\overline{G}$, then $G_<$ is a spanning subgraph of G.

Proof. Let G be a graph and $<$ an ordering of its vertices. As in the proof of Theorem 2, we will build a graph $G_<$ such that $V(G_<) = V(G)$, and $v < w$ are adjacent in $G_<$ if and only if they cannot belong to the same class of a partition which is strongly consistent with $<$. By definition of strong consistence, this happens if and only if either there is a vertex z in G such that $v < w < z$, z is adjacent to v and nonadjacent to w, or there is a vertex x such that $x < v < w$, x is adjacent to w and nonadjacent to v. So define $E(G_<)$ such that for $v < w$, $vw \in E(G_<)$ if and only if either there is a vertex z in G such that $v < w < z$, $zv \in E(G)$ and $zw \notin E(G)$ or there is a vertex x in G such that $x < v < w$, $xw \in E(G)$ and $xv \notin E(G)$.

Let us see that $<$ is a comparability ordering for $\overline{G_<}$. Suppose on the contrary that there is a triple $r < s < t$ in $V(G)$ such that rs, st are edges of $\overline{G_<}$ and rt is not an edge of $\overline{G_<}$. By definition of $G_<$, either there is a vertex z such that $r < s < t < z$, $zr \in E(G)$ and $zt \notin E(G)$, or there is a vertex x in G such that $x < r < s < t$, $xt \in E(G)$ and $xr \notin E(G)$. The case of z is identical to the one in the proof of Theorem 2. The case of x is symmetric, if $xs \notin E(G)$, then st is an edge of $G_<$, a contradiction. If $xs \in E(G)$, then rs is an edge of $G_<$, a contradiction as well. So $G_<$ is a co-comparability graph, being $<$ a comparability ordering for $\overline{G_<}$.

The relation with a coloring problem in $G_<$ is identical to the one in the proof of Theorem 2.

To complete the proof of the theorem, suppose now that G is a co-comparability graph and $<$ is a comparability ordering for $\overline{G}$. Let $v < w$ adjacent in $G_<$. By definition, either there is a vertex z in G such that $v < w < z$, $vz \in E(G)$ and $wz \notin E(G)$, or there is a vertex x in G such that $x < v < w$, $xw \in E(G)$ and $xv \notin E(G)$. Again, the case of z is identical to the one in the proof of Theorem 2, while the case of x is symmetric: if $vw \notin E(G)$, being $\overline{G}$ a comparability graph, $xw \notin E(G)$, a contradiction. So in any case $vw \in E(G)$. This proves that $G_<$ is a spanning subgraph of G. □

A direct consequence of this result is the following.

Corollary 13. *If G is a co-comparability graph,* $\mathrm{pthin}(G) \leq \chi(G)$. *Moreover, any vertex partition given by a coloring of G and any comparability ordering for its complement are strongly consistent.*

Again, in contrast with Theorem 12, if a partition is given, it is NP-complete to decide the existence of a strongly consistent ordering.

Strongly consistent ordering with a given partition
Instance: A graph $G = (V, E)$ and a partition of V into non-empty subsets.
Question: Does there exist a total order $<$ of V strongly consistent with the given partition?

By noticing that the ordering obtained from the solution to No-Betweenness instance in the proof of Theorem 5 is indeed strongly consistent with the vertex partition of the graph, we can prove the following.

Theorem 14. *The problem* Strongly consistent ordering with a given partition *is NP-complete.*

Also in this case, the computational complexity of the decision of existence of a strongly consistent ordering when the number of sets in the partition is fixed is still open. So is the computational complexity of deciding if a graph is proper k-thin, even for fixed $k \geq 2$, and even within the class of interval graphs.

3.3. Solving combinatorial optimization problems on graphs with bounded proper thinness

We start by the following observation: in a proper k-thin representation of a graph G, with ordering $<$ of V, namely $v_1 < \cdots < v_n$, and partition of V into k classes $V^1, \ldots, V^k$, for each pair of vertices $v_s < v_r$ that are in the same class, $N[v_s] \cap \{v_1, \ldots, v_s\} \supseteq N[v_r] \cap \{v_1, \ldots, v_s\}$. This allows us to handle other kinds of restrictions as for example domination type constraints.

Namely, if we are considering the subgraph G_s of G induced by $\{v_1, \ldots, v_s\}$ but we "keep in mind" that we still need to dominate some of the vertices in $\{v_{s+1}, \ldots, v_n\}$ with vertices of G_s, we can summarize these conditions into at

most k of them (each imposed by vertices of $\{v_{s+1}, \ldots, v_n\}$ in each partition class).

For graphs with bounded proper thinness k, when the proper k-thin representation of the graph is given, we can add now to the instance (with respect to Section 2.3) this kind of restrictions:

- $l_{ij(N)} \leq |S_i \cap N(v)| \leq u_{ij(N)} \quad \forall v \in S_j$, such that $l_{ij(N)} \in \{0, 1\}$ and $u_{ij(N)} \in \{1, \infty\}$ (it can be $i = j$), $1 \leq i, j \leq r$.
- $l_{ij[N]} \leq |S_i \cap N[v]| \leq u_{ij[N]} \quad \forall v \in S_j$, such that $l_{ij[N]} \in \{0, 1\}$ and $u_{ij[N]} \in \{1, \infty\}$ (it can be $i = j$), $1 \leq i, j \leq r$.

In this way the framework includes domination-type problems in the literature and their weighted versions, such as existence/minimum (weighted) independent dominating set, minimum (weighted) total dominating set, minimum perfect dominating set and existence/minimum (weighted) efficient dominating set, b-coloring [21] with fixed number of colors.

We will keep the notation of Section 2.3 and describe how to modify the algorithm in order to take into account the new restrictions. Now the vertex order and the partition of G are strongly consistent.

Each state now will be augmented with some new parameters:

- a family of nonnegative parameters $\{\gamma_{ij}\}_{1 \leq i \leq k; 1 \leq j \leq r}$, meaning that the last γ_{ij} vertices of V^i have already a neighbor in S_j (of index higher than them); they are kr and each of them may take a nonnegative value at most $n - 1$.
- a family of nonnegative parameters $\{\gamma^2_{ij}\}_{1 \leq i \leq k; 1 \leq j \leq r}$, meaning that the last γ^2_{ij} vertices of V^i have already two neighbors in S_j (of index higher than them); they are kr and each of them may take a nonnegative value at most $n - 1$.
- a family of nonnegative parameters $\{\lambda_{ijc}\}_{1 \leq i,c \leq k; 1 \leq j \leq r}$, meaning that, for each value $1 \leq c \leq k$, S_j has to contain at least one vertex in the set that is the union over $1 \leq i \leq k$ of the last λ_{ijc} vertices of V^i (if the union is empty, this mean no restriction associated with (c, S_j)); they are $k^2 r$ and each of them may take a nonnegative value at most $n - 1$.

The total number of states is then multiplied by at most n^{k^2r+2kr}, that keeps it polynomial in n, since k and r are constant.

The value of all these parameters in the only node of the layer X_n of the digraph is zero.

Now the problems associated with the nodes of X will have the additional constraints associated with the new restrictions and the parameters $\{\gamma_{ij}\}$, $\{\gamma_{ij}^2\}$, and $\{\lambda_{ijc}\}$.

Let us describe the additional conditions for the arcs of the digraph, whose labels and weights are still the same as in Section 2.3.

Let w be a node with parameters $(1, \ldots, \{\gamma_{ij}\}, \{\gamma_{ij}^2\}, \{\lambda_{ijc}\})$.

Let $1 \leq \ell \leq k$ such that $v_1 \in V^\ell$. For each $\tilde{J} \in L(v_1)$ (in particular $\tilde{J} \subseteq \{1, \ldots, r\}$) satisfying 1.1–1.5, and such that:

1.6 For each $1 \leq i \leq r$, $j \in \tilde{J}$, such that $l_{ij(N)} = 1$, $\gamma_{\ell i} > 0$.

1.7 For each $i \notin \tilde{J}$, $j \in \tilde{J}$, such that $l_{ij[N]} = 1$, $\gamma_{\ell i} > 0$.

1.8 For each $1 \leq i \leq r$, $j \in \tilde{J}$, such that $u_{ij(N)} = 1$ or $u_{ij[N]} = 1$, $\gamma_{\ell i}^2 = 0$.

1.9 For each $i, j \in \tilde{J}$, such that $u_{ij[N]} = 1$, $\gamma_{\ell i} = 0$.

1.10 For each $j \notin \tilde{J}$ and for each $1 \leq c \leq k$, $\lambda_{\ell jc} = 0$.

we add an arc from x_0 to w, labeled by $\tilde{J}$ and of weight $\sum_{1 \leq i \leq t; j \in \tilde{J}} c_{ij} w_i(v_1)$. If no $\tilde{J}$ satisfies conditions 1.1–1.10, no arc ending in w is added. If more than one arc x_0w was added, we can keep only the one with maximum (resp. minimum) weight if we are solving a maximization (resp. minimization) problem.

Note that if we add the arc x_0w labeled by $\tilde{J}$, then the solution $S_j = \{v_1\}$ for $j \in \tilde{J}$, $S_j = \emptyset$ for $j \notin \tilde{J}$ has weight $\sum_{1 \leq i \leq t; j \in \tilde{J}} c_{ij} w_i(v_1)$ and satisfies the state described by w: conditions 1.1–1.5 ensure the properties required in Section 2.3; conditions 1.6–1.9 ensure the validity of the two new families of restrictions about lower and upper bounds of neighbors of vertices of one set in other set, and condition 1.10 ensures that the restrictions imposed by the parameters $\{\lambda_{ijc}\}$ are satisfied.

Let w be a node with parameters $(s, \{l_{iJ\cap}\}, \{u_{iJ\cap}\}, \{l_{iJ\cup}\}, \{u_{iJ\cup}\}, \{\alpha_{ij}\}, \{\beta_{ij}\}, \{\gamma_{ij}\}, \{\gamma^2_{ij}\}, \{\lambda_{ijc}\})$, $1 < s \leq n$.

Let $1 \leq \ell \leq k$ such that $v_s \in V^\ell$. For each $\tilde{J} \in L(v_s)$ satisfying $s.1$–$s.4$, and such that:

$s.5$ For each $1 \leq i \leq r$, $j \in \tilde{J}$, such that $u_{ij(N)} = 1$ or $u_{ij[N]} = 1$, $\gamma^2_{\ell i} = 0$.

$s.6$ For each $i, j \in \tilde{J}$, such that $u_{ij[N]} = 1$, $\gamma_{\ell i} = 0$.

$s.7$ For each $j \notin \tilde{J}$ and for each $1 \leq c \leq k$, either $\lambda_{\ell jc} = 0$, or $\lambda_{\ell jc} > 1$, or there exists $1 \leq i \leq k$, $i \neq \ell$, such that $\lambda_{ijc} > 0$ (i.e., the union over $1 \leq i \leq k$ of the last λ_{ijc} vertices of V^i is not $\{v_s\}$).

$s.8$ For each $1 \leq i \leq r$ such that $\gamma_{\ell i} = 0$ and there exists $j \in \tilde{J}$ such that $l_{ij(N)} = 1$, $N(v_s) \cap \{1, \ldots, s-1\} \neq \emptyset$.

$s.9$ For each $i \notin \tilde{J}$ such that $\gamma_{\ell i} = 0$ and there exists $j \in \tilde{J}$ such that $l_{ij[N]} = 1$, $N(v_s) \cap \{1, \ldots, s-1\} \neq \emptyset$.

Let $\{\lambda^0_{ijc}\}_{1 \leq i,c \leq k; 1 \leq j \leq r}$ be defined this way: for every $j \in \tilde{J}$ and every $1 \leq c \leq k$ such that $\lambda_{\ell jc} > 0$, let $\lambda^0_{ijc} = 0$ for every $1 \leq i \leq k$; for every $j \in \tilde{J}$ and every $1 \leq c \leq k$ such that $\lambda_{\ell jc} = 0$, let $\lambda^0_{ijc} = \lambda_{ijc}$ for every $1 \leq i \leq k$; for every $j \notin \tilde{J}$ and every $1 \leq c \leq k$, let $\lambda^0_{\ell jc} = \max\{0, \lambda_{\ell jc} - 1\}$ and let $\lambda^0_{ijc} = \lambda_{ijc}$ for every $1 \leq i \leq k$, $i \neq \ell$.

Let $\{\lambda^1_{jc}\}_{1 \leq c \leq k; 1 \leq j \leq r}$ be defined as $\lambda^1_{jc} = 0$ if $\lambda^0_{ijc} = 0$ for every $1 \leq i \leq k$, $\lambda^1_{jc} = 1$ otherwise.

We add an arc from u to w, labeled by $\tilde{J}$ and of weight $\sum_{1 \leq i \leq t; j \in \tilde{J}} c_{ij} w_i(v_s)$, where u has parameters $(s-1, \{l'_{iJ\cap}\}, \{u'_{iJ\cap}\}, \{l'_{iJ\cup}\}, \{u'_{iJ\cup}\}, \{\alpha'_{ij}\}, \{\beta'_{ij}\}, \{\gamma'_{ij}\}, \{\gamma^{2'}{}_{ij}\}, \{\lambda'_{ijc}\})$, satisfies conditions $s'.2$–$s'.6$, and:

$s'.7$ For each $1 \leq i \leq r$ such that $\gamma_{\ell i} = 0$ and there exists $j \in \tilde{J}$ such that $l_{ij(N)} = 1$, if $\lambda^1_{i\ell} = 0$, then $\lambda'_{j'i\ell} = |N(v_s) \cap V^{j'} \cap \{1, \ldots, s-1\}|$ for each $1 \leq j' \leq k$; otherwise, $\lambda'_{j'i\ell} = \lambda^0_{j'i\ell}$ for every $1 \leq j' \leq k$ (recall that, by the observations above about proper thinness, $\lambda^0_{j'i\ell} = \min\{\lambda^0_{j'i\ell}, |N(v_s) \cap V^{j'} \cap \{1, \ldots, s-1\}|\}$).

$s'.8$ For each $i \notin \tilde{J}$ such that $\gamma_{\ell i} = 0$ and there exists $j \in \tilde{J}$ such that $l_{ij[N]} = 1$, if $\lambda^1_{i\ell} = 0$, then $\lambda'_{j'i\ell} = |N(v_s) \cap V^{j'} \cap \{1, \ldots, s-1\}|$ for each $1 \leq j' \leq k$; otherwise, $\lambda'_{j'i\ell} = \lambda^0_{j'i\ell}$ for every $1 \leq j' \leq k$.

$s'.9$ For each i, j, c not comprised in conditions $s'.7$ and $s'.8$, $\lambda'_{ijc} = \lambda^0_{ijc}$.

$s'.10$ Let $1 \leq j \leq r$. If there exists $j' \in \tilde{J}$ satisfying at least one of the following:

- $M_{jj'} = 0$
- $(u_{jj'(N)} = 1$ or $u_{jj'[N]} = 1)$ and $\gamma_{\ell j} > 0$
- $j \in \tilde{J}$ and $u_{jj'[N]} = 1$

then, $\beta'_{\ell j} = \max\{\beta_{\ell j} - 1, |N(v_s) \cap V^\ell \cap \{1, \ldots, s-1\}|\}$, and for $1 \leq i \leq k$, $i \neq \ell$, $\beta'_{ij} = \max\{\beta_{ij}, |N(v_s) \cap V^i \cap \{1, \ldots, s-1\}|\}$. Otherwise, $\beta'_{\ell j} = \max\{0, \beta_{\ell j} - 1\}$, and for $1 \leq i \leq k$, $i \neq \ell$, $\beta'_{ij} = \beta_{ij}$.

$s'.11$ For each $j \in \tilde{J}$: if $|N(v_s) \cap V^\ell \cap \{1, \ldots, s-1\}| \geq \gamma_{\ell j} - 1$, then $\gamma'_{\ell j} = |N(v_s) \cap V^\ell \cap \{1, \ldots, s-1\}|$ and $\gamma^{2'}_{\ell j} = \max\{0, \gamma_{\ell j} - 1\}$; otherwise, $\gamma'_{\ell j} = \max\{0, \gamma_{\ell j} - 1\}$ and $\gamma^{2'}_{\ell j} = \max\{\gamma^2_{\ell j} - 1, |N(v_s) \cap V^\ell \cap \{1, \ldots, s-1\}|\}$.

$s'.12$ For each $j \in \tilde{J}$, $1 \leq i \leq k$, $i \neq \ell$: if $|N(v_s) \cap V^i \cap \{1, \ldots, s-1\}| \geq \gamma_{ij}$, then $\gamma'_{ij} = |N(v_s) \cap V^i \cap \{1, \ldots, s-1\}|$ and $\gamma^{2'}_{ij} = \gamma_{ij}$; otherwise, $\gamma'_{ij} = \gamma_{ij}$ and $\gamma^{2'}_{ij} = \max\{\gamma^2_{ij}, |N(v_s) \cap V^i \cap \{1, \ldots, s-1\}|\}$.

If no $\tilde{J}$ satisfies conditions $s.1$–$s.9$, no arc ending in w is added. If more than one arc from the same vertex u to w was added, we can keep only the one with maximum (resp. minimum) weight if we are solving a maximization (resp. minimization) problem.

That is, if an arc is added, the arc corresponds to the choice of adding the vertex v_s to the sets $\{S_j\}_{j \in \tilde{J}}$, the conditions required imply that the choice is valid for w in the case that the state described by u admits a solution, the label of the arc keeps track of the choice made, and the cost corresponds to the weight that the choice adds to the objective function.

Note that if we add the arc uw labeled by $\tilde{J}$, then for a solution $\{S'_j\}_{1 \leq j \leq r}$ for G_{s-1} satisfying the state described by u, then the solution $\{S_j\}_{1 \leq j \leq r}$ for G_s such that $S_j = S'_j \cup \{v_s\}$ for $j \in \tilde{J}$, $S_j = S'_j$ for $j \notin \tilde{J}$ satisfies the state described by w.

Condition $s'.10$ ensures on one hand that the conditions imposed by the parameters $\{\beta_{ij}, u_{ij(N)}, u_{ij[N]}\}$ in w are satisfied by the solution of u, and, on the other hand, that if $j' \in \tilde{J}$ and $1 \leq j \leq r$ are such that $M_{jj'} = 0$ then no neighbor of v_s belongs to S'_j, as required. Conditions $s'.7$–$s'.9$ together with

$s.7$–$s.9$ define parameters $\{\lambda'_{ijc}\}$ in u in order to guarantee in w both the conditions imposed by the lower bounds $\{l_{ij(N)}, l_{ij[N]}\}$ and those imposed by the parameters $\{\lambda_{ijc}\}$. Finally, Condition $s'.11$ and $s'.12$ properly update the definition of parameters $\{\gamma'_{ij}, \gamma^{2'}{}_{ij}\}$ according to the choice $\tilde{J}$ for v_s. Condition $s'.2$–$s'.6$ are analyzed in Section 2.3.

As in that case, the difference of weight of the solution $\{S_j\}_{1\leq j\leq r}$ with respect to $\{S'_j\}_{1\leq j\leq r}$ is exactly $\sum_{1\leq i\leq t; j\in\tilde{J}} c_{ij} w_i(v_s)$.

In that way, a directed path in the digraph corresponds to an assignment of vertices of the graph to lists of sets and its weight is the value of the objective function for the corresponding assignment.

The digraph has a polynomial number of nodes and can be builded in polynomial time. Since it is acyclic, both the longest path and shortest path can be computed in linear time in the size of the digraph by topological sorting.

4. Thinness and graph operations

In this section we analyze the behavior of the thinness and proper thinness under different graph operations, namely union, join, and Cartesian product. The first two results allow us to fully characterize k-thin graphs by forbidden induced subgraphs within the class of cographs. The third result is used to solve in polynomial time the t-rainbow domination problem for fixed t on graphs with bounded proper thinness.

Let $G_1 = (V_1, E_1)$ and $G_2 = (V_2, E_2)$ be two graphs with $V_1 \cap V_2 = \emptyset$. The *union* of G_1 and G_2 is the graph $G_1 \cup G_2 = (V_1 \cup V_2, E_1 \cup E_2)$, and the *join* of G_1 and G_2 is the graph $G_1 \vee G_2 = (V_1 \cup V_2, E_1 \cup E_2 \cup V_1 \times V_2)$ (i.e., $\overline{G_1 \vee G_2} = \overline{G_1} \cup \overline{G_2}$).

Theorem 15. *Let G_1 and G_2 be graphs. Then* $\text{thin}(G_1 \cup G_2) = \max\{\text{thin}(G_1), \text{thin}(G_2)\}$ *and* $\text{pthin}(G_1 \cup G_2) = \max\{\text{pthin}(G_1), \text{pthin}(G_2)\}$.

Proof. Since both G_1 and G_2 are induced subgraphs of $G_1 \cup G_2$, then $\text{thin}(G_1 \cup G_2) \geq \max\{\text{thin}(G_1), \text{thin}(G_2)\}$ and the same holds for the proper thinness.

Let G_1 and G_2 be two graphs with thinness (resp. proper thinness) t_1 and t_2, respectively. Let $v_1, \ldots, v_{n_1}$ and $(V_1^1, \ldots, V_1^{t_1})$ be an ordering and

a partition of $V(G_1)$ which are consistent (resp. strongly consistent). Let $w_1, \ldots, w_{n_2}$ and $(V_2^1, \ldots, V_2^{t_2})$ be an ordering and a partition of $V(G_2)$ which are consistent (resp. strongly consistent). Suppose without loss of generality that $t_1 \leq t_2$. For $G = G_1 \cup G_2$, define a partition $V^1, \ldots, V^{t_2}$ such that $V^i = V_1^i \cup V_2^i$ for $i = 1, \ldots, t_1$ and $V^i = V_2^i$ for $i = t_1 + 1, \ldots, t_2$, and define $v_1, \ldots, v_{n_1}, w_1, \ldots, w_{n_2}$ as an ordering of the vertices. By definition of union of graphs, if three ordered vertices according to the order defined in $V(G_1 \cup G_2)$ are such that the first and the third are adjacent, either the three vertices belong to $V(G_1)$ or the three vertices belong to $V(G_2)$. Since the order and the partition restricted to each of G_1 and G_2 are the original ones, the properties required for consistence (resp. strong consistence) are satisfied. □

Theorem 16. *Let G_1 and G_2 be graphs. Then* $\text{thin}(G_1 \vee G_2) \leq \text{thin}(G_1) + \text{thin}(G_2)$ *and* $\text{pthin}(G_1 \vee G_2) \leq \text{pthin}(G_1) + \text{pthin}(G_2)$. *Moreover, if G_2 is complete, then* $\text{thin}(G_1 \vee G_2) = \text{thin}(G_1)$.

Proof. Let G_1 and G_2 be two graphs with thinness (resp. proper thinness) t_1 and t_2, respectively. Let $v_1, \ldots, v_{n_1}$ and $(V_1^1, \ldots, V_1^{t_1})$ be an ordering and a partition of $V(G_1)$ which are consistent (resp. strongly consistent). Let $w_1, \ldots, w_{n_2}$ and $(V_2^1, \ldots, V_2^{t_2})$ be an ordering and a partition of $V(G_2)$ which are consistent (resp. strongly consistent). For $G = G_1 \vee G_2$, define a partition with $t_1 + t_2$ sets as the union of the two partitions, and $v_1, \ldots, v_{n_1}, w_1, \ldots, w_{n_2}$ as an ordering of the vertices.

Let x, y, z be three vertices of $V(G)$ such that $x < y < z$, $xz \in E(G)$, and x and y are in the same class of the partition of $V(G)$. Then, in particular, x and y both belong either to $V(G_1)$ or to $V(G_2)$. If z belongs to the same graph, then $yz \in E(G)$ because the ordering and partition restricted to each of G_1 and G_2 are consistent. Otherwise, z is also adjacent to y by the definition of join.

We have proved that the defined partition and ordering are consistent, and thus that $\text{thin}(G_1 \vee G_2) \leq \text{thin}(G_1) + \text{thin}(G_2)$. The proof of the strong consistence, given the strong consistence of the partition and ordering of each of G_1 and G_2, is symmetric and implies $\text{pthin}(G_1 \vee G_2) \leq \text{pthin}(G_1) + \text{pthin}(G_2)$.

Suppose now that G_2 is complete (in particular, $t_2 = 1$). Since G_1 is an induced subgraph of $G_1 \vee G_2$, then $\text{thin}(G_1 \vee G_2) \geq \text{thin}(G_1)$. For $G =$

$G_1 \vee G_2$, define a partition $V^1, \ldots, V^{t_1}$ such that $V^1 = V_1^1 \cup V_2^1$ and $V^i = V_1^i$ for $i = 2, \ldots, t_1$, and define $v_1, \ldots, v_{n_1}, w_1, \ldots, w_{n_2}$ as an ordering of the vertices.

Let x, y, z be three vertices of $V(G)$ such that $x < y < z$, $xz \in E(G)$, and x and y are in the same class of the partition of $V(G)$. If z belongs to $V(G_2)$, then z is also adjacent to y, because it is adjacent to every vertex in $G - z$. If z belongs to $V(G_1)$, then x, y, and z, belong to $V(G_1)$ due to the definition of the order of the vertices, and thus $yz \in E(G)$ because the ordering and partition restricted to G_1 are consistent. This proves $\text{thin}(G_1 \vee G_2) \leq \text{thin}(G_1)$, and therefore $\text{thin}(G_1 \vee G_2) = \text{thin}(G_1)$. □

The following lemma shows a way of obtaining graphs with high thinness, using the join operator.

Lemma 17. *If G is not complete, then* $\text{thin}(G \vee 2K_1) = \text{thin}(G) + 1$.

Proof. By Theorem 16, $\text{thin}(G \vee 2K_1) \leq \text{thin}(G) + \text{thin}(2K_1) = \text{thin}(G) + 1$. On the other hand, as $G \vee 2K_1$ contains G as induced subgraph, $\text{thin}(G \vee 2K_1) \geq \text{thin}(G)$.

First notice that if $\text{thin}(G) = 1$ but G is not complete, then $G \vee 2K_1$ contains C_4 as induced subgraph, so it is not an interval graph, and $\text{thin}(G \vee 2K_1) \geq 2$, as claimed.

Suppose then that $\text{thin}(G) = k > 1$ and $\text{thin}(G \vee 2K_1) = k$ as well, and let $<$ be an ordering of the vertices of $G \vee 2K_1$ consistent with a partition $V^1, \ldots, V^k$. Let v, v' be the vertices of the graph $2K_1$, and suppose $v < v'$. Without loss of generality we may assume $v \in V^k$. As $\text{thin}(G) = k$, $V^k \cap V(G) \neq \emptyset$. Since $v' > v$, v' is nonadjacent to v, and v' is adjacent to all the vertices in $V^k \cap V(G)$, v has to be the smallest vertex in V^k. Let $z \in V^k \cap V(G)$ and suppose there is a vertex $x > z$ in $V(G)$. As x is adjacent to v', it is adjacent to z as well. So, we can define a new order $<'$ on $V(G \vee 2K_1)$ that preserves the order $<$ in $V^1 \cup V^{k-1} \cup \{v\}$ and such that the vertices of $V^k - \{v\}$ are the largest. By the observations above, this order $<'$ is still consistent with the partition $V^1, \ldots, V^k$. But it is also consistent with the partition $V^{1\prime}, \ldots, V^{k\prime}$ in which $V^{1\prime} = V^1 \cup V^k - \{v\}$, $V^{i\prime} = V^i$ for $1 < i < k$, and $V^{k\prime} = \{v\}$. This implies that $\text{thin}(G) < k$, a contradiction that completes the proof of the theorem. □

Cographs were defined in [9], where it was shown that they are exactly the graphs with no induced path of length four. Cographs admit a full decomposition theorem. Let the *trivial* graph be the one with one node only.

Proposition 18. [9] *Every cograph that is not trivial is either the union or the join of two smaller cographs.*

We will use this structural property along with the theorems about thinness of union and join of graphs to prove the following.

Theorem 19. *Let G be a cograph and $t \geq 1$. Then G has thinness at most t if and only if G contains no $\overline{(t+1)K_2}$ as induced subgraph.*

Proof. The only if part holds by Theorem 1, because the class of k-thin graph is hereditary for every k.

We will prove the if part by induction on the number of vertices of the cograph G. If G is a trivial graph, then $\text{thin}(G) = 1$ and the theorem holds. If G is not trivial, by Proposition 18, it is either union or join of two smaller cographs G_1 and G_2, with thinness t_1 and t_2, respectively.

Suppose first $G = G_1 \cup G_2$. By Theorem 15, $\text{thin}(G) = \max\{t_1, t_2\}$. If t_1 (resp. t_2) is greater than one, then by inductive hypothesis G_1 (resp. G_2) contains $\overline{t_1 K_2}$ (resp. $\overline{t_2 K_2}$) as induced subgraph, and so does G.

Suppose now that $G = G_1 \vee G_2$. If one of them is complete (suppose without loss of generality G_2), then, by Theorem 16, $\text{thin}(G) = t_1$. If t_1 is greater than one, then by inductive hypothesis G_1 contains $\overline{t_1 K_2}$ as induced subgraph, and so does G. If none of them is complete, then, by that fact and the inductive hypothesis, G_1 contains $\overline{t_1 K_2}$ and G_2 contains $\overline{t_2 K_2}$ as induced subgraph. As $\overline{t_1 K_2} \vee \overline{t_2 K_2} = \overline{(t_1 + t_2) K_2}$, G contains $\overline{(t_1 + t_2) K_2}$ as induced subgraph, thus $\text{thin}(G) \geq t1 + t2$ (Theorem 1). By Theorem 16, $\text{thin}(G) \leq t1 + t2$, and therefore $\text{thin}(G) = t1 + t2$. This finishes the proof of the theorem. □

A characterization by minimal forbidden induced subgraphs for k-thin graphs, $k \geq 2$, is open.

Let $G_1 = (V_1, E_1)$ and $G_2 = (V_2, E_2)$ be two graphs. The Cartesian product $G_1 \square G_2$ is a graph whose vertex set is the Cartesian product $V_1 \times V_2$, and such that two vertices (u_1, u_2) and (v_1, v_2) are adjacent in $G_1 \square G_2$ if and only if either $u_1 = v_1$ and u_2 is adjacent to v_2 in G_2, or $u_2 = v_2$ and u_1 is adjacent to v_1 in G_1.

Theorem 20. *Let G_1 and G_2 be graphs. Then* $\text{thin}(G_1 \Box G_2) \leq \text{thin}(G_1)|V(G_2)|$ *and* $\text{pthin}(G_1 \Box G_2) \leq \text{pthin}(G_1)|V(G_2)|$.

Proof. Let $G_1 = (V_1, E_1)$ be a k-thin (resp. proper k-thin) graph, and let $v_1, \ldots, v_{n_1}$ and $(V_1^1, \ldots, V_1^k)$ be an ordering and a partition of V_1 which are consistent (resp. strongly consistent). Let $G_2 = (V_2, E_2)$, $n_2 = |V_2|$, and $w_1, \ldots, w_{n_2}$ an arbitrary ordering of V_2. Consider $V_1 \times V_2$ lexicographically ordered with respect to the orderings of V_1 and V_2 above. Consider now the partition $\{V^{i,j}\}_{1 \leq i \leq k,\ 1 \leq j \leq n_2}$ such that $V^{i,j} = \{(v, w_j) : v \in V_1^i\}$ for each $1 \leq i \leq k$, $1 \leq j \leq n_2$. We will show that this ordering and partition of $V_1 \times V_2$ are consistent (resp. strongly consistent). Let $(v_p, w_i), (v_q, w_j), (v_r, w_\ell)$ be three vertices appearing in that ordering in $V_1 \times V_2$.

Case 1: $p = q = r$. In this case, the three vertices are in different classes, so no restriction has to be satisfied.

Case 2: $p = q < r$. In this case, (v_p, w_i) and (v_q, w_j) are in different classes. So suppose G_1 is proper k-thin and $(v_q, w_j), (v_r, w_\ell)$ belong to the same class, i.e., $j = \ell$. Vertices (v_p, w_i) and (v_r, w_ℓ) are adjacent in $G_1 \Box G_2$ if and only if $i = \ell$ and $v_p v_r \in E_1$. But then $(v_p, w_i) = (v_q, w_j)$, a contradiction.

Case 3: $p < q = r$. In this case, (v_q, w_j) and (v_r, w_ℓ) are in different classes. So suppose G_1 is k-thin and $(v_p, w_i), (v_q, w_j)$ belong to the same class, i.e., $i = j$. Vertices (v_p, w_i) and (v_r, w_ℓ) are adjacent in $G_1 \Box G_2$ if and only if $i = \ell$ and $v_p v_r \in E_1$. But then $(v_r, w_\ell) = (v_q, w_j)$, a contradiction.

Case 4: $p < q < r$. Suppose first G_1 is k-thin (resp. proper k-thin) and $(v_p, w_i), (v_q, w_j)$ belong to the same class, i.e., $i = j$ and v_p, v_q belong to the same class in G_1. Vertices (v_p, w_i) and (v_r, w_ℓ) are adjacent in $G_1 \Box G_2$ if and only if $i = \ell$ and $v_p v_r \in E_1$. But then $j = \ell$ and since the ordering and the partition are consistent (resp. strongly consistent) in G_1, $v_r v_q \in E_1$ and so (v_r, w_ℓ) and (v_q, w_j) are adjacent in $G_1 \Box G_2$. Now suppose that G_1 is proper k-thin and $(v_q, w_j), (v_r, w_\ell)$ belong to the same class, i.e., $j = \ell$. Vertices (v_p, w_i) and (v_r, w_ℓ) are adjacent in $G_1 \Box G_2$ if and only if $i = \ell$ and $v_p v_r \in E_1$. But then $i = j$ and since the ordering and the partition are strongly consistent in G_1, $v_p v_q \in E_1$ and so (v_p, w_i) and (v_q, w_j) are adjacent in $G_1 \Box G_2$. □

Corollary 21. *If G is proper k-thin then $G \Box K_t$ is proper kt-thin. In particular, if G is a proper interval graph then $G \Box K_t$ is proper t-thin.*

For a graph $G(V, E)$ and an integer t, we say that f is a *t-rainbow dominating function* if it assigns to each node $v \in V$ a subset of $\{1, \ldots, t\}$ such that $\cup_{u \in N(v)} f(u) = \{1, \ldots, t\}$ for all v with $f(v) = \emptyset$. Consider the following generalization of the dominating set problem.

t-RAINBOW DOMINATION PROBLEM
Instance: A graph $G = (V, E)$.
Find: a t-rainbow dominating function that minimizes $\sum_{v \in V} |f(v)|$.

The t-rainbow domination problem is equivalent to minimum dominating set of $G \Box K_t$ [6]. As a consequence of Corollary 21 and the results in Section 3.3, it can be solved in polynomial time on graphs with bounded proper thinness for fixed values of t. This generalizes the polynomiality for proper interval graphs, which was stated as an open question by Brešar and Kraner Šumenjak in [6]. The polynomiality of the t-rainbow domination problem for fixed t on interval graphs was recently proved by Hon, Kloks, Liu, and Wang in [20] (the algorithm for $t = 2$ is claimed in [19]).

5. Conclusions and open problems

We described a wide family of combinatorial optimization problems that can be solved in polynomial time on classes of bounded thinness and bounded proper thinness. We think that some restrictions can be further generalized (specially the domination type ones), with more involved sets of parameters and transition rules. We tried to keep it as simpler as possible, yet including many of the classical combinatorial optimization problems in the literature.

We also proved a number of theoretical results, some of them related with the recognition problem for the classes, others relating the concept of thinness and proper thinness with other known graph parameters, and analyzing their behavior under the graph operations union, join, and Cartesian product.

Some open problems are the following.

- Characterize (proper) k-thin graphs by minimal forbidden induced subgraphs (or at least within some graph class, we did it for thinness in cographs).
- What is the complexity of computing the thinness/proper thinness of a graph? Or deciding if it is at most k for some fixed values k?

- Can we develop some randomized algorithm to test just a subset of vertex orderings and obtain with high probability an approximation of the thinness/proper thinness?
- Can we improve the complexity of the algorithms to FPT? Or prove a hardness result?
- Given a partition of the vertex set into a *fixed* number k of classes, what is the complexity of deciding if there is a (strongly) consistent order for the vertices w.r.t. that partition (and finding it)? (We have proved that for an arbitrary number of classes the problems are NP-complete, and we have solved in polynomial time the symmetric problem, i.e., given the ordering, find a minimum (strongly) consistent partition.)
- Look at the relations with boolean width, cutwidth, and other width parameters (if there are any).
- Study the behavior of thinness under other graph products or graph operators in general.

Acknowledgements: This work was partially supported by UBACyT Grant 20020130100808BA, CONICET PIP 112-201201-00450CO, ANPCyT PICT 2012-1324 and 2015-2218 (Argentina). The authors want to thank Gianpaolo Oriolo for bringing our attention into the thinness parameter and for helpful discussions about it, Daniela Saban for the example of an interval graph with high proper thinness, Nicolas Nisse for pointing out the relation between proper thinness and other interval graphs invariants to us, Boštjan Brešar and Martin Milanič for pointing out the open problem on rainbow domination to us, and Yuri Faenza for his valuable help in the improvement of the manuscript.

References

[1] J. Arne Telle. Complexity of domination-type problems in graphs. *Nordic Journal of Computing*, 1(1):157–171, 1994.

[2] U. Bertelè and F. Brioschi. *Nonserial Dynamic Programming*. Academic Press, 1972.

[3] B.V.S. Bharadwaj and L.S. Chandran. Bounds on isoperimetric values of trees. *Discrete Mathematics*, 309(4):834–842, 2009.

[4] H.L. Bodlaender and K. Jansen. Restrictions of graph partition problems. Part I. *Theoretical Computer Science*, 148:93–109, 1995.

[5] F. Bonomo, S. Mattia, and G. Oriolo. Bounded coloring of cocomparability graphs and the pickup and delivery tour combination problem. *Theoretical Computer Science*, 412(45):6261–6268, 2011.

[6] B. Brešar and T. Kraner Šumenjak. On the 2-rainbow domination in graphs. *Discrete Applied Mathematics*, 155:2394–2400, 2007.

[7] M.R. Cerioli, F. de S. Oliveira, and J.L. Szwarcfiter. On counting interval lengths of interval graphs. *Discrete Applied Mathematics*, 159(7):532–543, 2011.

[8] S. Chandran, C. Mannino, and G. Oriolo. The indepedent set problem and the thinness of a graph. Manuscript, 2007.

[9] D. Corneil, H. Lerchs, and L. Stewart Burlingham. Complement reducible graphs. *Discrete Applied Mathematics*, 3(3):163–174, 1981.

[10] B. Courcelle, J. Engelfriet, and G. Rozenberg. Handle-rewriting hypergraph grammars. *Journal of Computer and System Sciences*, 46(2):218–270, 1993.

[11] B. Courcelle and S. Olariu. Upper bounds to the clique width of graphs. *Discrete Applied Mathematics*, 101(1–3):77–114, 2000.

[12] X. Deng, P. Hell, and J. Huang. Linear time representation algorithms for proper circular-arc graphs and proper interval graphs. *SIAM Journal on Computing*, 25(2):390–403, 1996.

[13] J.A. Ellis, I.H. Sudborough, and J.S. Turner. The vertex separation and search number of a graph. *Information and Computation*, 113(1):50–79, 1994.

[14] T. Feder, P. Hell, S. Klein, L.T. Nogueira, and F. Protti. List matrix partitions of chordal graphs. *Theoretical Computer Science*, 349(1):52–66, 2005.

[15] P.C. Fishburn. *Interval orders and interval graphs: A study of partially ordered sets*. John Wiley & Sons, 1985.

[16] M.C. Golumbic. The complexity of comparability graph recognition and coloring. *Computing*, 18:199–208, 1977.

[17] M.C. Golumbic and U. Rotics. On the clique-width of some perfect graph classes. *International Journal of Foundations of Computer Science*, 11(3):423–443, 2000.

[18] W. Guttmann and M. Maucher. Variations on an ordering theme with constraints. In G. Navarro, L. Bertossi, and Y. Kohayakawa, editors, *Proceedings of the Fourth IFIP International Conference on Theoretical Computer Science - TCS 2006*, volume 209 of *IFIP International Federation for Information Processing*, pages 77–90. Springer US, 2006.

[19] W-K. Hon, T. Kloks, H-H. Liu, and H-L. Wang. Rainbow domination and related problems on some classes of perfect graphs. In M. Hajiaghayi and M. Mousavi, editors, *Topics in Theoretical Computer Science - TTCS 2015*, volume 9541 of *Lecture Notes in Computer Science*, pages 121–134. Springer, Cham, 2016.

[20] W.-K. Hon, T. Kloks, H.-H. Liu, and H.-L. Wang. Rainbow domination and related problems on some classes of perfect graphs. arXiv:1502.07492v1 [cs.DM], February 2015.

[21] R. W. Irving and D. F. Manlove. The b-chromatic number of a graph. *Discrete Applied Mathematics*, 91:127–141, 1999.

[22] K. Jansen. The optimum cost chromatic partition problem. In G. Bongiovanni, D.P. Bovet, and G. Di Battista, editors, *Proceedings of the Italian Conference on Algorithms and Complexity - CIAC 1997*, volume 1203 of *Lecture Notes in Computer Science*, pages 25–36, 1997.

[23] P. Klavík, Y. Otachi, and J. Šejnoha. On the classes of interval graphs of limited nesting and count of lengths. In Seok-Hee Hong, editor, *27th International Symposium on Algorithms and Computation (ISAAC 2016)*, volume 64 of *Leibniz International Proceedings in Informatics (LIPIcs)*, pages 45:1–45:13, Dagstuhl, Germany, 2016.

[24] R. Leibowitz, S.F. Assmann, and G.W. Peck. The interval count of a graph. *SIAM Journal on Algebraic and Discrete Methods*, 3:485–494, 1982.

[25] Z. Lonc. On complexity of some chain and antichain partition problems. In G. Schmidt and R. Berghammer, editors, *Proceedings of the International Workshop on Graph-Theoretic Concepts in Computer Science 1991*, volume 570 of *Lecture Notes in Computer Science*, pages 97–104. Springer, Berlin, Heidelberg, 1992.

[26] P. Looges and S. Olariu. Optimal greedy algorithms for indifference graphs. *Computers & Mathematics with Applications*, 25:15–25, 1993.

[27] C. Mannino, G. Oriolo, F. Ricci, and S. Chandran. The stable set problem and the thinness of a graph. *Operations Research Letters*, 35:1–9, 2007.

[28] N. Nisse, 2010. Personal communication.

[29] S. Olariu. An optimal greedy heuristic to color interval graphs. *Information Processing Letters*, 37:21–25, 1991.

[30] E. Prisner. Graphs with few cliques. In Y. Alavi and A. Schwenk, editors, *Proceedings of the 7th Quadrennial International Conference on the Theory and Applications of Graphs, 1992*, Graph Theory, Combinatorics, and Applications, pages 945–956. Wiley Interscience, New York, 1995.

[31] G. Ramalingam and C. Pandu Rangan. A unified approach to domination problems on interval graphs. *Information Processing Letters*, 27:271–274, 1988.

[32] F.S. Roberts. Indifference graphs. In F. Harary, editor, *Proof Techniques in Graph Theory*, pages 139–146. Academic Press, 1969.

[33] F.S. Roberts. On the boxicity and cubicity of a graph. In W.T. Tutte, editor, *Recent Progress in Combinatorics*, pages 301–310. Academic Press, 1969.

[34] N. Robertson and P.D. Seymour. Graph minors. I. Excluding a forest. *Journal of Combinatorial Theory. Series B*, 35(1):39–61, 1983.

[35] N. Robertson and P.D. Seymour. Graph minors. II. Algorithmic aspects of tree-width. *Journal of Algorithms*, 7(3):309–322, 1986.

[36] D. Saban, 2010. Personal communication.

[37] P. Scheffler. A linear algorithm for the pathwidth of trees. In R. Bodendiek and R. Henn, editors, *Topics in Combinatorics and Graph Theory: Essays in Honour of Gerhard Ringel*, pages 613–620. Physica-Verlag HD, Heidelberg, 1990.

[38] S. Tsukiyama, M. Idle, H. Ariyoshi, and Y. Shirakawa. A new algorithm for generating all the maximal independent sets. *SIAM Journal on Computing*, 6(3):505–517, 1977.

[39] Zs. Tuza. Graph colorings with local constraints – a survey. *Discussiones Mathematicae. Graph Theory*, 17:161–228, 1997.

[40] I. Vrt'o. A note on isoperimetric peaks of complete trees. *Discrete Mathematics*, 310(6–7):1272–1274, 2010.

[41] D. West. *Introduction to Graph Theory*. Prentice Hall, 2nd edition, 2000.